

Research article

Open Access

Effects of β -alanine supplementation and high-intensity interval training on endurance performance and body composition in men; a double-blind trial

Abbie E Smith¹, Ashley A Walter², Jennifer L Graef¹, Kristina L Kendall¹, Jordan R Moon¹, Christopher M Lockwood¹, David H Fukuda¹, Travis W Beck², Joel T Cramer² and Jeffrey R Stout*¹

Address: ¹Metabolic and Body Composition Laboratory, Department of Health and Exercise Science, University of Oklahoma, Norman, OK 73019, USA and ²Biophysics Laboratory; Department of Health and Exercise Science, University of Oklahoma, Norman, OK 73019, USA

Email: Abbie E Smith - abbiesmith@ou.edu; Ashley A Walter - ashannwalter@ou.edu; Jennifer L Graef - jennifer-l-graef-1@ou.edu; Kristina L Kendall - krissykendall@ou.edu; Jordan R Moon - jordanmoon@ou.edu; Christopher M Lockwood - clockwood@muscleandfitness.com; David H Fukuda - david.fukuda@ou.edu; Travis W Beck - tbeck@ou.edu; Joel T Cramer - jrcramer@ou.edu; Jeffrey R Stout* - jrstout@ou.edu

* Corresponding author

Published: 11 February 2009

Received: 8 January 2009

Journal of the International Society of Sports Nutrition 2009, **6**:5 doi:10.1186/1550-2783-6-5

Accepted: 11 February 2009

This article is available from: <http://www.jissn.com/content/6/1/5>

© 2009 Smith et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Intermittent bouts of high-intensity exercise result in diminished stores of energy substrates, followed by an accumulation of metabolites, promoting chronic physiological adaptations. In addition, β -alanine has been accepted has an effective physiological hydrogen ion (H^+) buffer. Concurrent high-intensity interval training (HIIT) and β -alanine supplementation may result in greater adaptations than HIIT alone. The purpose of the current study was to evaluate the effects of combining β -alanine supplementation with high-intensity interval training (HIIT) on endurance performance and aerobic metabolism in recreationally active college-aged men.

Methods: Forty-six men (Age: 22.2 ± 2.7 yrs; Ht: 178.1 ± 7.4 cm; Wt: 78.7 ± 11.9 ; VO_{2peak} : 3.3 ± 0.59 l·min⁻¹) were assessed for peak O_2 utilization (VO_{2peak}), time to fatigue (VO_{2TTE}), ventilatory threshold (VT), and total work done at 110% of pre-training VO_{2peak} (TWD). In a double-blind fashion, all subjects were randomly assigned into one either a placebo (PL – 16.5 g dextrose powder per packet; n = 18) or β -alanine (BA – 1.5 g β -alanine plus 15 g dextrose powder per packet; n = 18) group. All subjects supplemented four times per day (total of 6 g/day) for the first 21-days, followed by two times per day (3 g/day) for the subsequent 21 days, and engaged in a total of six weeks of HIIT training consisting of 5–6 bouts of a 2:1 minute cycling work to rest ratio.

Results: Significant improvements in VO_{2peak} , VO_{2TTE} , and TWD after three weeks of training were displayed ($p < 0.05$). Increases in VO_{2peak} , VO_{2TTE} , TWD and lean body mass were only significant for the BA group after the second three weeks of training.

Conclusion: The use of HIIT to induce significant aerobic improvements is effective and efficient. Chronic BA supplementation may further enhance HIIT, improving endurance performance and lean body mass.

Background

High-intensity exercise results in diminished stores of adenosine tri-phosphate (ATP), phosphocreatine (PCr) and glycogenic substrates, and the intracellular accumulation of metabolites (adenosine di-phosphate (ADP), inorganic phosphate (P_i), hydrogen ions (H^+) and magnesium (Mg^+), each of which has been implicated as a cause of muscle fatigue [1-3]. Excessive formation of H^+ results in a decrease in intramuscular pH which may contribute to fatigue in some models of exercise [1,4-6]. Enhancing an individual's ability to buffer protons may delay fatigue by improving the use of energy substrates and maintaining muscular contraction [6-9]. When the time and intensity level of exercise is sufficient, the majority of protons that are produced are buffered by the bicarbonate (HCO_3^-) buffering system [10,11] in which they are exported from the muscle [12]. Physiological buffering during dynamic exercise is typically controlled by the HCO_3^- system and is also supported by direct physico-chemical buffering, provided mainly by phosphate, histidine residues of peptides and proteins, and the small amount of bicarbonate present in muscle at the start of exercise. However, during short bursts of intense exercise, such as HIIT, physico-chemical buffering will exceed that by HCO_3^- mediated dynamic buffering, calling on intramuscular stores of phosphates and peptides.

Specifically, carnosine (β -alanyl-L-histidine), a cytoplasmic dipeptide, constitutes an important non-bicarbonate physico-chemical buffer. By virtue of a pKa of 6.83 and its high concentration in muscle, carnosine is more effective at sequestering protons than either bicarbonate (pKa 6.37) or inorganic phosphate (pKa 7.2), the other two major physico-chemical buffers over the physiological pH range [7,13]. However, as a result of the greater concentration of carnosine in muscle than bicarbonate in the initial stages of muscle contraction, and inorganic phosphate, its buffering contribution may be quantitatively more important.

Mechanisms for increasing muscle carnosine concentration have been somewhat disputed. While carnosine may be increased in chronically trained athletes, the effects of acute training are less clear. In one study, it has been reported that eight weeks of intensive training may increase intramuscular carnosine content [14]. In contrast, several other studies have shown that intense training, of up to 16 weeks, has been unable to promote a rise in skeletal muscle carnosine levels [6,15-17]. Only when β -alanine supplementation was combined with training did an increase in muscle carnosine occur [16], although the increase (40–60%) was similar to that seen with supplementation alone [18].

While carnosine is synthesized in the muscle from its two constituents, β -alanine and histidine [19], synthesis is

limited by the availability of β -alanine [18,20]. β -alanine supplementation alone has been shown to significantly increase the intramuscular carnosine content [6,18]. Elevation of intramuscular carnosine content via β -alanine supplementation alone, has been shown to improve performance [6,14,21-24]. Recently, Hill and colleagues [6] demonstrated a 13% improvement in total work done (TWD) following four weeks of β -alanine supplementation, and an additional 3.2% increase after 10 weeks. Zoeller et al. [24] also reported significant increases in ventilatory threshold (VT) in a sample of untrained men after supplementing with β -alanine ($3.2 \text{ g} \cdot \text{d}^{-1}$) for 28 days. In agreement, Kim et al. [21] also reported significant increases in VT and time to exhaustion (TTE) in highly trained male cyclists after 12 weeks of β -alanine ($4.8 \text{ g} \cdot \text{d}^{-1}$) supplementation and endurance training. Furthermore, Stout et al. [22,23] reported a significant delay in neuromuscular fatigue, measured by physical working capacity at the fatigue threshold (PWC_{FT}), in both men and women after 28 days of β -alanine supplementation ($3.2 \text{ g} \cdot \text{d}^{-1} - 6.4 \text{ g} \cdot \text{d}^{-1}$). Despite the improvements in VT, TTE, TWD, and PWC_{FT} after supplementation, there were no increases in aerobic power, measured by VO_{2peak} [22-24].

Although HIIT alone does not appear to increase skeletal muscle carnosine content [17], training has been suggested to improve muscle buffering capacity [25-27]. When repeated bouts of high-intensity intervals are interspersed with short rest periods, subsequent trials are initiated at a much lower pH [28]. Training in such a manner subjects the body to an acidic environment, forcing several physiological adaptations. Notably, HIIT has been shown to improve VO_{2peak} and whole body fat oxidation in only two weeks (7 sessions at 90% VO_{2peak}) [29]. Furthermore, over a longer period of time (4–6 weeks), HIIT has been reported to increase high-intensity exercise performance (6–21%), muscle buffering capacity, whole body exercise fat oxidation, and aerobic power (VO_{2peak}) [25-27].

The respective supporting bodies of literature for the use of β -alanine supplementation alone and high-intensity training alone have gained recent popularity. However, to date, no study has combined and evaluated concurrent HIIT with β -alanine supplementation. In theory, we hypothesize that an increase in intramuscular carnosine content, as a result of β -alanine supplementation, may enhance the quality of HIIT by reducing the accumulation of hydrogen ions, leading to greater physiological adaptations. Therefore, the purpose of this study was to determine the effects of chronic (6 weeks) β -alanine supplementation in combination with HIIT on endurance performance measures in recreationally trained individuals.

Methods

Subjects

Forty-six college-aged men, who were recreationally active one to five hours per week, and had not taken any sports supplement within the six months prior-, volunteered to participate in this study (mean \pm SD; Age: 22.2 ± 2.7 yrs, Height: 178.1 ± 7.4 cm, Weight: 78.7 ± 11.9 kg). Subjects were informed of the potential risks, benefits, and time requirements prior to enrolling and giving written consent. All study procedures were approved by the University's Institutional Review Board.

Study design

This double-blind, randomized study included two three-week periods of HIIT and β -alanine supplementation. All participants completed a series of baseline, mid- and post-testing, including a series of cycling tests and body composition assessment using air displacement plethysmography (BodPod[®]) at all time points. Following baseline testing subjects were randomly assigned, in a double-blind fashion, to one of two supplementing groups, β -alanine or placebo, both with HIIT. Participant's initial VO_{2peak} power output values were used to establish the TWD intensity and the training intensity for the six week duration, with no modification to intensity following mid-testing. The first three-week period of training was completed at workloads between 90%–110% of each individual's VO_{2peak} , while the second three-week training peaked at 115%. While training, participants supplemented with 6 g per day of β -alanine or placebo during the first three weeks and 3 g per day during the second three week phase. Supplementing with 6.4 g per day of β -alanine, for 28 days has demonstrated a 60% increase in carnosine concentration [6,18], supporting the 21 day phase, allowing for an adequate loading period for β -alanine to elicit increases in intramuscular carnosine concentration. Furthermore, recent literature suggests even greater increases in carnosine levels when combining high-intensity training and β -alanine supplementation [17]. Following the three-week adaptation phase, mid-training and post-training tests were completed in the same order as the pre-testing, allowing at least 48 hours between each testing session. All subjects were instructed to maintain their current diet throughout the duration of the study and were asked to refrain from caffeine and vigorous activity 24 hours prior to any testing session. Food logs were distributed to all participants and completed (two non-consecutive weekdays and one weekend day) at baseline-testing, mid-testing and post-testing, to evaluate any changes in total kcal and/or protein intake.

Determination of VO_{2peak}

At pre-, mid-, and post-training, all participants performed a continuous graded exercise test (GXT) on an electronically braked cycle ergometer (Corval 400, Gonin-

gen, The Netherlands) to determine VO_{2peak} , time to exhaustion (VO_{2TTE}) and ventilatory threshold (VT). Pedal cadence was maintained at 70 rpm, while the power output was initially set at 50 W for a five minute warm-up, and increased by 25 W every two minutes, until the participant could no longer maintain the required power output (cadence dropped below 60 rpm). Respiratory gases were monitored breath by breath and analyzed with open-circuit spirometry (True One 2400[®] Metabolic Measurement System, Parvo-Medics Inc., Provo UT) to determine VO_{2peak} and VT. The data was averaged over 15 second intervals. The highest 15 second VO_2 value during the GXT was recorded as the VO_{2peak} value if it coincided with at least two of the following criteria: (a) a plateau in heart rate (HR) or HR values within 10% of the age-predicted HRmax, (b) a plateau in VO_2 (defined by an increase of note more than $150 \text{ ml} \cdot \text{min}^{-1}$), and/or (c) an RER value greater than 1.15 [30]. Heart rate was also monitored continuously during exercise by using a heart rate monitor (Polar FS1, Polar Electro Inc. Lake Success, NY). The amount of time to reach exhaustion (VO_{2TTE}) during the VO_{2peak} was also recorded in seconds. Ventilatory threshold (VT) was determined using standard software (True One 2400[®] Metabolic Measurement System, Parvo-Medics Inc., Provo UT) by plotting ventilation (V_E) against VO_2 as described previously [31]. Two linear regression lines were fit to the lower and upper portions of the V_E vs. VO_2 curve, before and after the break points, respectively. The intersection of these two lines was defined as VT, and was recorded with respect to the corresponding power output (W).

Test-retest reliability for the VO_{2peak} protocol at the University of Oklahoma using twenty-one men, demonstrate reliable between-day testing with an intraclass correlation coefficient (ICC) of 0.975 (SEM $0.257 \text{ l} \cdot \text{min}^{-1}$) and a percent of coefficient of variation (%CV) of 5.18%.

Total Work Done Cycling Test

Each subject performed a constant-load time to exhaustion (TTE) test on an electronically braked cycle ergometer, at a cadence of ~ 70 rpm. Participants performed a five minute warm-up at 50 W, followed by a cycle to exhaustion at their individual pre-determined workload, established at 110% of the maximum VO_{2peak} workload (W). The subject's TTE was defined by the time (in seconds), that could be maintained without dropping below a cadence of 60 rpm. Total work done (TWD) was further calculated as the primary variable of interest, using the product of time (in seconds) and the power output (W), divided by 1,000, and presented in kilojoules (kJ).

The reliability statistics for TWD reflect a strong ICC of 0.713 (SEM 25.2 kJ) and a %CV of 3.80%.

Training intervention and -alanine supplementation

Training was performed on an electronically braked cycle ergometer (Corval 400, Groningen, The Netherlands) to maintain testing specificity. Participants began the supervised training session within two to four days following testing. Following the baseline-testing and group randomization, subjects began the first of two, three-week training periods. Training followed a fractal periodized plan to allow for adequate progression and to prevent overtraining [32] and was completed three days per week. The training intensity began at 90% of the maximum power output (W) achieved during the baseline VO₂ peak test and progressed in an undulating manner, reaching a maximum of 115% by the end of the second, three-week training period. The first three-week period consisted of five sets of two-minute intervals with one-minute rest periods. The second three-week session followed a similar protocol, modifying the progression by increasing the repetitions from five to six, during weeks six and seven and still taking place on three days per week (Figure 1). A training log was completed for each training session. The total time (seconds) completed and workload (watts) was used to compute total training volume (kJ) (Figure 2).

alanine (1.5 g β-alanine, 15 g dextrose per dose) or placebo (16.5 g dextrose per dose). Supplements were mixed with water in an orange flavored dextrose powder and were consumed four times throughout the day. On the three days that subjects visited the lab for training, they consumed two pre-mixed doses, one 30 minutes before, and one immediately after completion of the training session. The remaining two doses were taken that day, *ad libitum*. For the remaining four days of the week, participants were instructed to mix and consume the four doses (6 g per day) of their respective supplement, *ad libitum*. Throughout the second three-week training period, participants supplemented in a similar manner for on- and off-training days, for an additional 21 days, at a dose of 3 g per day, taken in two, 16.5 g doses (1.5 g β-alanine, 15 g dextrose). The participants in the placebo group consumed an isovolumetric flavored powder (16.5 g dextrose) identical in appearance and taste to the β-alanine. Participants were asked to record each dose on a designated dosing log for each day and they were asked to bring in the supplement packaging to allow investigators to monitor compliance.

In addition to training, during the first three-week period, the participants also supplemented with 6 g per day β-

Determination of body composition

Body composition was assessed prior-to, mid-way, and following training and supplementing by using air dis-




Figure 1
Training protocol for the first and second three-week training phases, respectively. Black represents five sets of the 2:1 training, while grey represents six sets of the same 2:1 protocol.




Figure 2
2A. The average \pm SD weekly training load (2A; watts) and training time (2B; seconds) between the BA (black) and PL (grey) treatment groups, across the six-week training protocol.

placement plethysmography (Bod Pod®). The subjects' weight (kg) and body volume were measured and used to determine percent body fat, fat mass (kg), and lean body mass (kg) using the revised formula of Brozek et al. [33].

Statistical analysis

Separate two-way repeated measures ANOVAs (group [β -alanine vs. placebo] \times time [pre- vs. mid- vs. post-supplementation]) were used to identify any group by time interactions. If a significant interaction occurred, the statistical model was decomposed by examining the simple main effects with separate one-way repeated measures ANOVAs for each group and one-way factorial ANOVAs for each time. An alpha of $p \leq 0.05$ was used to determine statistical significance. All data are reported as mean \pm standard deviation (SD).

Results

Table 1 presents the mean and standard deviation values for VO_{2peak} ($l \cdot min^{-1}$), VO_{2TTE} (seconds), VT (watts) and TWD (kJ) for both treatment groups at pre-, mid- and post-testing.

VO_{2peak} , VO_{2TTE} , VT during GXT

Significant main effects for time resulted for maximal oxygen consumption (VO_{2peak}), time to exhaustion (VO_{2TTE}) and ventilatory threshold (VT) determined during the graded exercise ($p < 0.001$). There were significant improvements in VO_{2peak} after three weeks of training and supplementing across both treatment groups ($p < 0.001$; ES: 0.977). While there were no significant difference for the improvements in VO_{2peak} at any time point between groups, only the BA group demonstrated significant improvements from mid- to post-training and supplementing ($p = 0.010$) with no significant change from mid- to post- for the PL group ($p = 0.118$). Similar results for VO_{2TTE} were also revealed with both groups demonstrating significant improvements from pre- to mid-testing ($p < 0.001$; ES: 0.983), with no difference between groups. Significant changes from mid- to post- VO_{2TTE} were only evident in the BA group ($p = 0.043$).

There were no significant differences among the improvements in VT between groups. Improvements from pre- to mid VT for both the PL and BA groups did not yield significance. However, the PL group was the only group to dem-

Table 1: Mean \pm SD values for VO_{2peak} (l·min⁻¹), VO_{2TTE} (s), VT (W) and TWD (kJ) at pre-, mid-, and post-testing.

		Maximal Oxygen Consumption (l·min ⁻¹)		Time to Exhaustion (s)		Ventilatory Threshold (W)		Total Work Done (kJ)	
		β -alanine	Placebo	β -alanine	Placebo	β -alanine	Placebo	β -alanine	Placebo
Pre-test	Mean	3.28	3.25	1168.2	1128.7	140.3	127.3	58.4	55.7
	SD	0.57	0.63	163.6	166.9	35.5	42.6	19.2	13.8
Mid-test	Mean	3.52*	3.56*	1304.9*	1258.7*	154.2	140.3	89.0*	83.3*
	SD	0.49	0.56	153.7	204.5	36.6	52.3	30.1	25.7
Post-test	Mean	3.67†	3.66	1386.7†	1299.6	172.2	188.9†	131.3†	102.0†
	SD	0.58	0.55	234.9	164.9	65.2	58.3	81.7	36.7

*indicates a significant difference from pre- to mid-testing ($p < 0.05$)

†indicates a significant improvement from mid- to post-testing ($p < 0.05$)

onstrate significant improvements from mid- to post ($p = 0.001$).

Time to exhaustion test-TWD

The improvements in TWD were significant across all time points, with no difference between groups ($p > 0.05$; ES: 0.898). While not significant, the delta values at both time points were greater for the BA group [pre-mid: 30.6 ± 19.9 sec; mid-post: 42.3 ± 72.1 sec] when compared to the PL group [pre-mid: 27.6 ± 22.1 ; mid-post: 18.6 ± 28.3].

Body Composition

The physical characteristics of the subjects determined at mid-testing and after six-weeks of HIIT and supplementing are presented in Table 2. Body mass did not change significantly with supplementing or training. However, the determination of body composition with the use of air displacement plethysmography (Bod Pod[®]) revealed a significant improvement from pre- to mid-testing in lean body mass in only the BA group ($p = 0.011$; ES: 0.985) and no change in the PL group ($p = 0.138$). Furthermore, there were no significant changes in percent body fat ($p = 0.287$) or fat mass ($p = 0.984$) between treatment groups after three and six weeks of HIIT and supplementing.

Dietary Analysis

There was no significant difference between groups for their supplement or training compliance rate, representing a 6.4 -3.2 g per day intake for the BA group, for the three and six weeks, respectively. Analyses of the dietary

recalls demonstrated no significant differences in caloric intake ($p > 0.05$) between the BA (3120 ± 244 kcal) and placebo (2775 ± 209 kcal) groups. Furthermore, there were no differences in macronutrient daily intake, with both groups consuming 47% of their daily calories from carbohydrates, 34% from fat and 16% from protein.

Training Volume

There was a significant main effect for time ($p < 0.01$) for both training volume (watts) and training time (seconds). However, there was no significant difference between groups for either volume (Figure 2A) or time (Figure 2B), at any time point (weeks 1–6). Although not significant, the BA group consistently trained at higher workloads and for longer time periods.

Discussion

The current study is the first to examine the effects of concurrent high-intensity interval training (HIIT) and β -alanine supplementation on a series of physiological and performance variables. The primary findings support the use of HIIT as an advantageous training tool. Furthermore, the current study also proposes the use of β -alanine supplementation to enhance the benefits of HIIT, by possibly improving muscle buffer capacity after six weeks of training and supplementing. The maximal oxygen uptake and time to reach maximum oxygen consumption (VO_{2peak} , VO_{2TTE}) and total work done (TWD) increased significantly in both training groups (β -alanine and placebo) over a six week HIIT protocol (Table 1). However,

Table 2: Mean \pm SD values for body weight (kg), body fat (%), lean body mass (kg), and fat mass (kg) from pre-, mid-, and post-testing.

	β -alanine (n = 18)			Placebo (n = 18)		
	Pre-testing	Mid-testing	Post-testing	Pre-testing	Mid-testing	Post-testing
Weight (kg)	78.8 \pm 12.8	80.1 \pm 13.0	79.8 \pm 12.4	78.5 \pm 11.3	79.3 \pm 12.3	79.8 \pm 11.9
Body Fat (%)	13.7 \pm 6.3	13.7 \pm 6.4	13.7 \pm 5.6	16.1 \pm 7.5	15.9 \pm 8.3	16.0 \pm 7.9
Lean Body Mass (kg)	67.6 \pm 8.9	68.6 \pm 8.6*	68.4 \pm 8.4	65.5 \pm 8.1	66.1 \pm 8.5	65.8 \pm 8.4
Fat Mass (kg)	11.3 \pm 6.5	11.5 \pm 6.8	11.3 \pm 6.0	13.0 \pm 7.1	13.1 \pm 8.0	13.0 \pm 7.8

*indicates a significant difference from pre- to mid-testing. ($p < 0.05$).

β -alanine supplementation appeared to have a greater influence on $\text{VO}_{2\text{peak}}$ and $\text{VO}_{2\text{TTE}}$, resulting in a significant ($p < 0.05$) increase during the second three weeks of training, while no change occurred in placebo group. In addition, TWD significantly ($p < 0.05$) increased during the last three weeks by 32% and 18% for the β -alanine and Placebo groups, respectively. Improvements in VT were also reported for both training groups, however the placebo group demonstrated significant improvements during the last three week training phase (Table 1). Lastly, the present study also identified a significant change in lean body mass for the β -alanine supplementing group after three weeks, with no change in the placebo group.

Enhanced $\text{VO}_{2\text{peak}}$, $\text{VO}_{2\text{TTE}}$, and VT after training

A series of HIIT interventions have suggested that interval exercise ($> 80\% \text{VO}_{2\text{max}}$) elicits greater gains in aerobic capacity than moderate-intensity exercise [34-36]. Consequently, the improvements reported in cardiorespiratory fitness in the current study were similar to most studies that have employed short-term (2-9 weeks) endurance interval training programs in untrained and recreationally active individuals [25,29,34,37-40]. Specifically, the average reported increases in $\text{VO}_{2\text{peak}}$ have ranged from 6-20% in male and female populations. Although the training regimens utilized have varied slightly, all supporting studies applied a similar protocol. The use of a 1:1 [37,38,40] and a 2:1 [29,34,39] work-to-rest design (1-4 minutes) have been the most effective for promoting an increase in aerobic capacity. Our data supports previous literature, suggesting a 7-10% increase in $\text{VO}_{2\text{peak}}$ during the first three week training phase and a 3-4.5% increase following the second three week session. While both groups significantly improved in $\text{VO}_{2\text{peak}}$ and $\text{VO}_{2\text{TTE}}$ from pre- to mid-testing, only the β -alanine group demonstrated significant improvements from mid- to post-testing (Table 1).

The use of high-intensity exercise as a training modality has been shown to stimulate acute and chronic physiological adaptations (cardiovascular, metabolic, respiratory and neural), which ultimately lead to improved performance [34,37,41]. The increases in $\text{VO}_{2\text{peak}}$, $\text{VO}_{2\text{TTE}}$, and VT reported in the current study are in line with other studies, which have suggested that the improvements in aerobic performance are attributable to a reduction in anaerobic ATP production, resulting from an increased contribution of aerobic energy production at higher intensity workloads [42,43]. The greater reliance on aerobic metabolism for energy has been further linked to an up-regulation of various glycolytic enzymes (phosphofruktokinase, hexokinase, citrate synthetase, and sodium potassium ATPase) [42,44-47], as well as with increased mitochondrial density and improved blood flow due to increased capillarization [44,45]. These improvements, in

combination with an enhanced ability to buffer H^+ , may provide some explanation into the greater improvements in the second three-week training phase, in the BA group only. Although blood pH levels were not measured directly, support from training volume (Figure 2A) and training time (Figure 2B), demonstrate that participants supplementing with β -alanine engaged in longer, more intense training sessions, possibly leading to greater adaptations.

Improvements in TWD

In addition to augmenting $\text{VO}_{2\text{peak}}$, $\text{VO}_{2\text{TTE}}$ and VT, the HIIT program utilized in the current study demonstrated significant improvements in TWD (Table 1). Interestingly, the increases in total work performed in the current study were greater than in previously reported improvements in TWD following HIIT alone [48-50], with both groups demonstrating a 50-53% improvement during the first three weeks of training and the β -alanine group showing a 32% increase compared to the 18% increase in the placebo group, after the second three-week training phase. In support, Kim et al. [21] demonstrated significantly greater increases in TWD in highly trained cyclists after a 12-week β -alanine supplementation and endurance training program, compared to training only. In addition, Hill et al. [6] also demonstrated significant improvements in TWD (13%) on a cycle ergometer following four weeks of β -alanine supplementation, without training. While the data appear to support the use of β -alanine supplementation to augment TWD, with and without training, the previously mentioned studies utilized highly trained participants, compared to an un-trained population in the current study.

Scientists have suggested the use of β -alanine may enhance training adaptations [6,18,23], by increasing ability to train at a higher intensity without fatigue. Recently Harris et al. [18] and Hill et al. [6] have posited that increasing skeletal muscle carnosine concentration with β -alanine supplementation may improve the ability to stabilize the intramuscular pH during intense exercise by buffering accumulating H^+ . Offsetting the indirect effect of proton accumulation on contractile function with the use of β -alanine, has been shown to be effective in delaying neuromuscular fatigue, improving VT and time to exhaustion in both trained and untrained individuals [6,21,23,24]. Furthermore, Kim et al. [21] reported a significant increase in VT after 12 weeks of endurance and resistance training while supplementing β -alanine in highly trained cyclists. However, our results demonstrated no added benefit of combining β -alanine supplementation and HIIT to elicit increases in VT, greater than training alone. The differences in training status (elite vs. recreationally trained) may have resulted in the conflicting results between the current study and Kim and col-

leagues. Additional research examining the effects of concurrent β -alanine supplementation and HIIT in trained versus untrained men and women would provide additional insight toward the current findings.

Augmented Lean Body Mass

Interestingly, the improvements in performance over the six-weeks of training also demonstrated concomitant gains in lean body mass in the β -alanine group only. Recent evidence suggests that intense exercise may elicit intramuscular acidosis, potentially augmenting protein degradation [51], inhibiting protein synthesis [52] and thus hindering training adaptations. Another theory posited suggests that β -alanine supplementation may have allowed for greater training volume thus providing a greater stimulus, resulting in significant gains in lean body mass, as observed in the current study. In support, Hoffman et al. [53,54] reported significantly higher training volume for athletes consuming β -alanine during resistance training sessions, which they hypothesized lead to significant increases in lean body mass. In short, minimizing the acidic response from HIIT, and/or increasing training volume with β -alanine supplementation, may help to increase lean body mass and lead to improvements in performance.

Conclusion

Our findings support the use of HIIT as an effective training stimulus for improving aerobic performance, in as little as three weeks. The use of β -alanine supplementation, in combination with HIIT, appeared to result in greater changes in VO_2 peak and VO_{2TTE} during the second three weeks of training, while no significant change occurred in placebo group. In addition, TWD significantly ($p < 0.05$) increased during the last three weeks by 32% and 18% for the β -alanine and Placebo groups, respectively. While more research is needed, the current study suggests that in untrained young men, the use of β -alanine supplementation may enhance the benefits of HIIT and augment endurance performance.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to this work. All authors have read and approved the final manuscript.

Acknowledgements

A sincere thanks is given to Dr. Roger Harris, University of Chichester, Chichester, UK, for his time and input he contributed to reviewing this manuscript.

The authors would also like to thank FSI Nutrition, 2132 South 156th Circle, Omaha, NE <http://www.fsinutrition.com> and RunFast Promotions, 8790

Wendy Lane South, West Palm Beach FL, 33411 <http://www.runfastpromotions.com> for supporting and funding this research endeavor.

References

1. Robergs RA, Ghiasvand F, Parker D: **Biochemistry of exercise-induced metabolic acidosis.** *Am J Physiol Regul Integr Comp Physiol* 2004, **287(3)**:R502-516.
2. Spriet LL, Lindinger MI, McKelvie RS, Heigenhauser GJ, Jones NL: **Muscle glycogenolysis and H⁺ concentration during maximal intermittent cycling.** *Journal of applied physiology* 1989, **66(1)**:8-13.
3. Allen DG, Lamb GD, Westerblad H: **Skeletal muscle fatigue: cellular mechanisms.** *Physiological reviews* 2008, **88(1)**:287-332.
4. Messonnier L, Kristensen M, Juel C, Denis C: **Importance of pH regulation and lactate/H⁺ transport capacity for work production during supramaximal exercise in humans.** *J Appl Physiol* 2007, **102(5)**:1936-1944.
5. Pottenger JA, Webster MJ, Nickel GL, Haub MD, Palmer RJ: **The effects of buffer ingestion on metabolic factors related to distance running performance.** *European journal of applied physiology and occupational physiology* 1996, **72(4)**:365-371.
6. Hill CA, Harris RC, Kim HJ, Harris BD, Sale C, Boobis LH, Kim CK, Wise JA: **Influence of beta-alanine supplementation on skeletal muscle carnosine concentrations and high intensity cycling capacity.** *Amino acids* 2007, **32(2)**:225-233.
7. Abe H: **Role of histidine-related compounds as intracellular proton buffering constituents in vertebrate muscle.** *Biochemistry* 2000, **65(7)**:757-765.
8. Suzuki Y, Nakao T, Maemura H, Sato M, Kamahara K, Morimatsu F, Takamatsu K: **Carnosine and anserine ingestion enhances contribution of nonbicarbonate buffering.** *Medicine and science in sports and exercise* 2006, **38(2)**:334-338.
9. McNaughton L, Backx K, Palmer G, Strange N: **Effects of chronic bicarbonate ingestion on the performance of high-intensity work.** *European journal of applied physiology and occupational physiology* 1999, **80(4)**:333-336.
10. Beaver WL, Wasserman K, Whipp BJ: **Bicarbonate buffering of lactic acid generated during exercise.** *J Appl Physiol* 1986, **60(2)**:472-478.
11. Juel C: **Regulation of pH in human skeletal muscle: adaptations to physical activity.** *Acta physiologica (Oxford, England)* 2008, **193(1)**:17-24.
12. Juel C, Klarskov C, Nielsen JJ, Krstrup P, Mohr M, Bangsbo J: **Effect of high-intensity intermittent training on lactate and H⁺ release from human skeletal muscle.** *American journal of physiology* 2004, **286(2)**:E245-251.
13. Bate-Smith EC: **The buffering of muscle in rigor: Protein, phosphate and carnosine.** *The Journal of physiology* 1938, **92**:336-343.
14. Suzuki Y, Ito O, Mukai N, Takahashi H, Takamatsu K: **High level of skeletal muscle carnosine contributes to the latter half of exercise performance during 30-s maximal cycle ergometer sprinting.** *The Japanese journal of physiology* 2002, **52(2)**:199-205.
15. Derave W, Ozdemir MS, Harris RC, Pottier A, Reyngoudt H, Koppo K, Wise JA, Achten E: **beta-Alanine supplementation augments muscle carnosine content and attenuates fatigue during repeated isokinetic contraction bouts in trained sprinters.** *J Appl Physiol* 2007, **103(5)**:1736-1743.
16. Harris RC, Edge J, Kendrick IP, Bishop D, Goodman C, Wise JA: **The Effect of Very High Interval Training on the Carnosine Content and Buffering Capacity of V Lateralis from Humans.** *FASEB J* 2007, **21**:769.
17. Kendrick IP, Harris RC, Kim HJ, Kim CK, Dang VH, Lam TQ, Bui TT, Smith M, Wise JA: **The effects of 10 weeks of resistance training combined with beta-alanine supplementation on whole body strength, force production, muscular endurance and body composition.** *Amino acids* 2008, **34(4)**:547-554.
18. Harris RC, Tallon MJ, Dunnett M, Boobis L, Coakley J, Kim HJ, Fallowfield JL, Hill CA, Sale C, Wise JA: **The absorption of orally supplied beta-alanine and its effect on muscle carnosine synthesis in human vastus lateralis.** *Amino acids* 2006, **30(3)**:279-289.
19. Bakardjiev A, Bauer K: **Transport of beta-alanine and biosynthesis of carnosine by skeletal muscle cells in primary culture.** *European journal of biochemistry/FEBS* 1994, **225(2)**:617-623.
20. Dunnett M, Harris RC, Soliman MZ, Suwar AA: **Carnosine, anserine and taurine contents in individual fibres from the middle**

- gluteal muscle of the camel. *Research in veterinary science* 1997, **62(3)**:213-216.
21. Kim HJ, Kim CK, Lee YW, Harris RC, Sale C, Harris BD, Wise JA: **The effect of a supplement containing B-alanine on muscle carnosine synthesis and exercise capacity, during 12 week combined endurance and weight training.** *J Int Soc Sports Nutr* 2006, **3**:S9.
 22. Stout JR, Cramer JT, Mielke M, O'Kroy J, Torok DJ, Zoeller RF: **Effects of twenty-eight days of beta-alanine and creatine monohydrate supplementation on the physical working capacity at neuromuscular fatigue threshold.** *Journal of strength and conditioning research/National Strength & Conditioning Association* 2006, **20(4)**:928-931.
 23. Stout JR, Cramer JT, Zoeller RF, Torok D, Costa P, Hoffman JR, Harris RC, O'Kroy J: **Effects of beta-alanine supplementation on the onset of neuromuscular fatigue and ventilatory threshold in women.** *Amino acids* 2007, **32(3)**:381-386.
 24. Zoeller RF, Stout JR, O'Kroy JA, Torok DJ, Mielke M: **Effects of 28 days of beta-alanine and creatine monohydrate supplementation on aerobic power, ventilatory and lactate thresholds, and time to exhaustion.** *Amino acids* 2007, **33(3)**:505-510.
 25. Edge J, Bishop D, Goodman C: **The effects of training intensity on muscle buffer capacity in females.** *European journal of applied physiology* 2006, **96(1)**:97-105.
 26. Laursen PB, Jenkins DG: **The scientific basis for high-intensity interval training: optimising training programmes and maximising performance in highly trained endurance athletes.** *Sports medicine (Auckland, NZ)* 2002, **32(1)**:53-73.
 27. Weston AR, Myburgh KH, Lindsay FH, Dennis SC, Noakes TD, Hawley JA: **Skeletal muscle buffering capacity and endurance performance after high-intensity interval training by well-trained cyclists.** *European journal of applied physiology and occupational physiology* 1997, **75(1)**:7-13.
 28. Costill DL, Verstaappen F, Kuipers H, Janssen E, Fink W: **Acid-base balance during repeated bouts of exercise: influence of HCO₃.** *International journal of sports medicine* 1984, **5(5)**:228-231.
 29. Talanian JL, Galloway SD, Heigenhauser GJ, Bonen A, Spriet LL: **Two weeks of high-intensity aerobic interval training increases the capacity for fat oxidation during exercise in women.** *Journal of applied physiology* 2007, **102(4)**:1439-1447.
 30. Day JR, Rossiter HB, Coats EM, Skasick A, Whipp BJ: **The maximally attainable VO₂ during exercise in humans: the peak vs. maximum issue.** *J Appl Physiol* 2003, **95(5)**:1901-1907.
 31. Orr GW, Green HJ, Hughson RL, Bennett GW: **A computer linear regression model to determine ventilatory anaerobic threshold.** *J Appl Physiol* 1982, **52(5)**:1349-1352.
 32. Brown L, Greenwood M: **Periodization Essentials and Innovations in Resistance Training Protocols.** *JSCR* 2005, **27(4)**:80-85.
 33. Brozek J, Grande F, Anderson JT, Keys A: **Densitometric Analysis of Body Composition: Revision of Some Quantitative Assumptions.** *Annals of the New York Academy of Sciences* 1963, **110**:113-140.
 34. Helgerud J, Hoydal K, Wang E, Karlsen T, Berg P, Bjerkaas M, Simonsen T, Helgesen C, Hjørth N, Bach R, et al.: **Aerobic high-intensity intervals improve VO₂max more than moderate training.** *Medicine and science in sports and exercise* 2007, **39(4)**:665-671.
 35. Rognmo O, Hetland E, Helgerud J, Hoff J, Stordahl SA: **High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease.** *Eur J Cardiovasc Prev Rehabil* 2004, **11(3)**:216-222.
 36. Thomas TR, Adeniran SB, Etheridge GL: **Effects of different running programs on VO₂ max, percent fat, and plasma lipids.** *Canadian journal of applied sport sciences* 1984, **9(2)**:55-62.
 37. Berger NJ, Tolfrey K, Williams AG, Jones AM: **Influence of continuous and interval training on oxygen uptake on-kinetics.** *Medicine and science in sports and exercise* 2006, **38(3)**:504-512.
 38. Burke J, Thayer R, Belcamino M: **Comparison of effects of two interval-training programmes on lactate and ventilatory thresholds.** *British journal of sports medicine* 1994, **28(1)**:18-21.
 39. Duffield R, Edge J, Bishop D: **Effects of high-intensity interval training on the VO₂ response during severe exercise.** *Journal of science and medicine in sport/Sports Medicine Australia* 2006, **9(3)**:249-255.
 40. Eddy DO, Sparks KL, Adelizi DA: **The effects of continuous and interval training in women and men.** *European journal of applied physiology and occupational physiology* 1977, **37(2)**:83-92.
 41. Rozenek R, Funato K, Kubo J, Hoshikawa M, Matsuo A: **Physiological responses to interval training sessions at velocities associated with VO₂max.** *Journal of strength and conditioning research/National Strength & Conditioning Association* 2007, **21(1)**:188-192.
 42. Gaitanos GC, Williams C, Boobis LH, Brooks S: **Human muscle metabolism during intermittent maximal exercise.** *Journal of applied physiology* 1993, **75(2)**:712-719.
 43. Harmer AR, McKenna MJ, Sutton JR, Snow RJ, Ruell PA, Booth J, Thompson MW, Mackay NA, Stathis CG, Crameri RM, et al.: **Skeletal muscle metabolic and ionic adaptations during intense exercise following sprint training in humans.** *Journal of applied physiology* 2000, **89(5)**:1793-1803.
 44. Henriksson J: **Effects of physical training on the metabolism of skeletal muscle.** *Diabetes care* 1992, **15(11)**:1701-1711.
 45. Krusturup P, Hellsten Y, Bangsbo J: **Intense interval training enhances human skeletal muscle oxygen uptake in the initial phase of dynamic exercise at high but not at low intensities.** *The Journal of physiology* 2004, **559(Pt 1)**:335-345.
 46. Nordborg N, Bangsbo J, Pilegaard H: **Effect of high-intensity training on exercise-induced gene expression specific to ion homeostasis and metabolism.** *Journal of applied physiology* 2003, **95(3)**:1201-1206.
 47. Rodas G, Ventura JL, Cadefau JA, Cusso R, Parra J: **A short training programme for the rapid improvement of both aerobic and anaerobic metabolism.** *European journal of applied physiology* 2000, **82(5-6)**:480-486.
 48. Coggan AR, Kohrt WM, Spina RJ, Kirwan JP, Bier DM, Holloszy JO: **Plasma glucose kinetics during exercise in subjects with high and low lactate thresholds.** *Journal of applied physiology* 1992, **73(5)**:1873-1880.
 49. Demarle AP, Heugas AM, Slawinski JJ, Tricot VM, Koralsztein JP, Billat VL: **Whichever the initial training status, any increase in velocity at lactate threshold appears as a major factor in improved time to exhaustion at the same severe velocity after training.** *Archives of physiology and biochemistry* 2003, **111(2)**:167-176.
 50. Gaiga MC, Docherty D: **The effect of an aerobic interval training program on intermittent anaerobic performance.** *Canadian journal of applied physiology = Revue canadienne de physiologie appliquee* 1995, **20(4)**:452-464.
 51. Caso G, Garlick PJ: **Control of muscle protein kinetics by acid-base balance.** *Current opinion in clinical nutrition and metabolic care* 2005, **8(1)**:73-76.
 52. Ballmer PE, Imoberdorf R: **Influence of acidosis on protein metabolism.** *Nutrition (Burbank, Los Angeles County, Calif)* 1995, **11(5)**:462-468. discussion 470.
 53. Hoffman J, Ratamess N, Faigenbaum A, Ross R, Kang J, Stout J, Wise JA: **Short-duration beta-alanine supplementation increases training volume and reduces subjective feelings of fatigue in college football players.** *Nutrition Research* 2007, **28(1)**:31-35.
 54. Hoffman J, Ratamess N, Kang J, Mangine G, Faigenbaum A, Stout J: **Effect of creatine and beta-alanine supplementation on performance and endocrine responses in strength/power athletes.** *International journal of sport nutrition and exercise metabolism* 2006, **16(4)**:430-446.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

